

**BILKENT UNIVERSITY**  
Mathematics Department

**Math 116 Intermediate Calculus III**  
**Summer School 2006-2007**

**FIRST MIDTERM EXAM - Answer Key**

---

---

*4:00 pm - 6:00 pm (120 minutes)*

*June 17, 2007*

---

**Surname** : .....

**Name** : .....

**Id. No.** : .....

**Section** : .....

---

**IMPORTANT**

- This exam consists of 5 questions of equal weight.
- Each question is on a separate sheet. Please read the questions carefully and write your answers under the corresponding question. Be neat.
- Show all your work. Correct answers without sufficient explanation might not get the full credit.
- Calculators are not allowed.

Please do not write anything below this line.

---

Q1	Q2	Q3	Q4	Q5	Total
20 pts.	20 pts.	20 pts.	20 pts.	20 pts.	100 pts.

**Question 1** (10+10=20 points). Let

$$f(x, y) = \begin{cases} \frac{x^3 \cos x - 4y^3}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

(a) Show that  $f$  is continuous at  $(0, 0)$ .

Note that

$$\left| \frac{x^3 \cos x - 4y^3}{x^2 + y^2} - 0 \right| \leq \frac{|x|x^2 + 4|y|y^2}{x^2 + y^2} \leq |x| + 4|y| \leq 5\sqrt{x^2 + y^2}$$

Thus, for any given  $\epsilon > 0$ , choose  $\delta > 0$  as  $\delta < \epsilon/5$ , so that

$$\left| \frac{x^3 \cos x - 4y^3}{x^2 + y^2} - 0 \right| < \epsilon \quad \text{when} \quad 0 < \sqrt{x^2 + y^2} < \delta.$$

This proves that  $f(x, y)$  is continuous at  $(0, 0)$  since  $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = 0$ .

(b) Compute  $f_x(0, 0)$  and  $f_y(0, 0)$  by using definition of partial derivatives.

$f(0, 0) = 0$ ,  $f(h, 0) = h \cos(h)$  and  $f(0, k) = -4k$ . Then

$$f_x(0, 0) = \lim_{h \rightarrow 0} \frac{f(h, 0) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{h \cos(h)}{h} = 1,$$

and

$$f_y(0, 0) = \lim_{k \rightarrow 0} \frac{f(0, k) - f(0, 0)}{k} = \lim_{k \rightarrow 0} \frac{-4k}{k} = -4.$$

**Question 2** (10+10=20 points).

(a) Let  $u(x, y) = xyf\left(\frac{x+y}{xy}\right)$  where  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a differentiable function. Show that

$$x^2 \frac{\partial u}{\partial x} - y^2 \frac{\partial u}{\partial y} = G(x, y)u(x, y),$$

for some function  $G(x, y)$  of two variables and find  $G(x, y)$ .

It just follows from the Chain rule:

$$\frac{\partial u}{\partial x} = yf\left(\frac{x+y}{xy}\right) + xy \frac{xy - y(x+y)}{x^2y^2} f'\left(\frac{x+y}{xy}\right)$$

Therefore,

$$\frac{\partial u}{\partial x} = yf\left(\frac{x+y}{xy}\right) - \frac{y}{x} f'\left(\frac{x+y}{xy}\right) \quad \text{and} \quad \frac{\partial u}{\partial y} = xf\left(\frac{x+y}{xy}\right) - \frac{x}{y} f'\left(\frac{x+y}{xy}\right)$$

Finally,

$$x^2 \frac{\partial u}{\partial x} - y^2 \frac{\partial u}{\partial y} = (x-y)xyf\left(\frac{x+y}{xy}\right) \quad \text{and hence,} \quad G(x, y) = x - y.$$

(b) Let  $f(x, y, z) = e^x \cos(yz)$ . Find the directional derivative of  $f$  at the origin in the direction of  $\mathbf{v} = 3\mathbf{i} + 6\mathbf{j} - 2\mathbf{k}$ .

The unit vector  $\mathbf{u}$  in direction of  $\mathbf{v}$  is given by

$$\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{3}{7}\mathbf{i} + \frac{6}{7}\mathbf{j} - \frac{2}{7}\mathbf{k}$$

Since  $\nabla f = e^x \cos(yz)\mathbf{i} - ze^x \sin(yz)\mathbf{j} - ye^x \sin(yz)\mathbf{k}$ . Then  $\nabla f|_{(0,0,0)} = \mathbf{i}$  and

$$\left(D_{\mathbf{u}}f\right)_{(0,0,0)} = \nabla f|_{(0,0,0)} \cdot \mathbf{u} = 3/7.$$

**Question 3** (10+10=20 points). Let  $f(x, y) = \frac{1}{2}x^2 + xy + \frac{1}{4}y^2 + 3\cos(x - 2) - 3y + 4$ .

(a) Find the linearization  $L(x, y)$  of  $f(x, y)$  at the point  $P(2, 1)$ .

We have  $f_x(x, y) = x + y - 3\sin(x - 2)$  and  $f_y(x, y) = x + y/2 - 3$ . So,  $f_x(2, 1) = 3$  and  $f_y(2, 1) = -1/2$  also  $f(2, 1) = 33/4$ .

Thus the linearization  $L(x, y)$  of  $f(x, y)$  at the point  $P(2, 1)$  is given by

$$L(x, y) = \frac{33}{4} + 3(x - 2) - \frac{1}{2}(y - 1)$$

(b) Find an upper bound for the magnitude  $|E(x, y)|$  of the error term  $E(x, y)$  in the standard linear approximation  $f(x, y) \approx L(x, y)$  over the rectangle  $R : |x - 2| \leq 0.1$  and  $|y - 1| \leq 0.3$

Now, we have  $f_{xx}(x, y) = 1 - 3\cos(x - 2)$   $f_{xy}(x, y) = 1$  and  $f_{yy}(x, y) = 1/2$ . Hence,  $|f_{xx}(x, y)| \leq 4$ ,  $|f_{xy}(x, y)| \leq 4$  and  $|f_{yy}(x, y)| \leq 4$ .

Since,

$$|E(x, y)| \leq \frac{M}{2} (|x - 2| + |y - 1|)^2$$

We can take  $M = 4$  and  $|x - 2| \leq 0.1$  and  $|y - 1| \leq 0.3$ , then  $|E(x, y)| \leq 0.32$ .

**Question 4** (20 points). The temperature function at all points in the disc  $x^2 + y^2 \leq 1$  is given by  $T(x, y) = (x + y)e^{-(x^2 + y^2)}$ . Find the highest and lowest temperatures at the points of the disc. Give the coordinates of points for the highest and lowest temperatures.

$$T_x(x, y) = (1 - 2x^2 - 2xy)e^{-(x^2 + y^2)} \text{ and } T_y(x, y) = (1 - 2y^2 - 2xy)e^{-(x^2 + y^2)}$$

The critical points are the solutions of the system below:

$$\begin{cases} 1 - 2x^2 - 2xy = 0 \\ 1 - 2y^2 - 2xy = 0 \end{cases}$$

So,  $y = x$  or  $y = -x$ . The former implies that  $1 - 2x^2 = 2x^2$ , i.e.  $x = \pm 1/2$  and the latter implies  $1 + 2x^2 = 2x^2$  and consequently, there is no solution for the case of  $y = -x$ . Therefore, the critical points are  $(1/2, 1/2)$  and  $(-1/2, -1/2)$  and also note that  $T(1/2, 1/2) = 1/\sqrt{e}$  and  $T(-1/2, -1/2) = -1/\sqrt{e}$ .

Use the method of Lagrange's Multipliers to find extremum values of the temperature function  $T(x, y)$  on the boundary  $x^2 + y^2 = 1$ , i.e.  $\nabla T = \lambda \nabla g$ , where  $g(x, y) = x^2 + y^2 - 1 = 0$ . More precisely,

$$(1 - 2x^2 - 2xy)e^{-(x^2 + y^2)}\mathbf{i} + (1 - 2y^2 - 2xy)e^{-(x^2 + y^2)}\mathbf{j} = \lambda(2x\mathbf{i} + 2y\mathbf{j}).$$

Since  $x$  and  $y$  are nonzero,

$$\frac{x}{y} = \frac{1 - 2x^2 - 2xy}{1 - 2y^2 - 2xy},$$

then  $x = y$  and  $x^2 + y^2 = 1$ , i.e.  $(1/\sqrt{2}, 1/\sqrt{2})$  and  $(-1/\sqrt{2}, -1/\sqrt{2})$  are the critical points of  $T(x, y)$  on the boundary.

Notice that  $T(1/\sqrt{2}, 1/\sqrt{2}) = \sqrt{2}/e$  and  $T(-1/\sqrt{2}, -1/\sqrt{2}) = -\sqrt{2}/e$ .

Therefore, the highest temperature  $1/\sqrt{e}$  is taken at  $(1/2, 1/2)$ , and the lowest temperature  $-1/\sqrt{e}$  is taken at  $(-1/2, -1/2)$ .

**Question 5** (4+8+8=20 points). Let  $f(x, y) = e^{x^2-4y} + \ln(x - y^2)$  and  $P(2, 1)$  be given.

(a) Find an equation of the level curve  $C$  of  $f$  that passes through  $P(2, 1)$ .

$f(2, 1) = e^0 + \ln 1 = 1$ , so the level curve  $C$  is

$$e^{x^2-4y} + \ln(x - y^2) = 1.$$

(b) Find an equation of the tangent line to the level curve  $C$  in part (a) at the point  $P(2, 1)$ .

We have,

$$\nabla f = \left( 2xe^{x^2-4y} + \frac{1}{x-y^2} \right) \mathbf{i} - \left( 4e^{x^2-4y} + \frac{2y}{x-y^2} \right) \mathbf{j}$$

and

$$\nabla f|_{(2,1)} = 5\mathbf{i} - 6\mathbf{j}.$$

then the equation of the tangent line is

$$5(x - 2) - 6(y - 1) = 0.$$

or  $5x - 6y = 4$

(c) Find an equation of the tangent plane of the surface  $z = f(x, y)$  at the point  $(2, 1, f(2, 1))$ .

Define  $g(x, y, z) = z - f(x, y)$ , then

$$\nabla g = -\left( 2xe^{x^2-4y} + \frac{1}{x-y^2} \right) \mathbf{i} + \left( 4e^{x^2-4y} + \frac{2y}{x-y^2} \right) \mathbf{j} + \mathbf{k},$$

and hence  $\nabla g|_{(2,1,f(2,1))} = -5\mathbf{i} + 6\mathbf{j} + \mathbf{k}$ .

The tangent plane to the surface  $z = f(x, y)$  at the point  $(2, 1, f(2, 1))$  is given by

$$-5(x - 2) + 6(y - 1) + (z - 1) = 0.$$

or  $5x - 6y - z = 3$