

FIRST PROJECT OF MATH544: ON CALCULUS OF

VARIATIONS *February, 2013.*

Prove all theorems and lemmas in the text.

In the theory of extrema of functions of single variable, a necessary condition for a minimum, besides $f'(x) = 0$, is that $f'' \geq 0$ (if it exists). A condition somewhat analogous to this holds for functionals. Let us suppose that there is an admissible function y , for which $J(y) = \int_a^b L(x, y, y')dx$ is a minimum. Then $f(\epsilon) = J(y + \epsilon h)$ has a minimum at $\epsilon = 0$; accordingly $f'(0) = 0$ (from which follows the Euler-Lagrange equation $E_y(L) = 0$) and $f''(0) \geq 0$, assuming its existence. Hence for all h ,

$$f''(0) = \int_a^b [L_{yy} h^2 + 2L_{yy'} h h' + L_{y'y'} h'^2]dx \geq 0$$

Choose a special variation

$$h = \begin{cases} 0 & a \leq x \leq \xi - \epsilon \\ 1 + (x - \xi)/\epsilon & \xi - \epsilon \leq x \leq \xi \\ 1 - (x - \xi)/\epsilon & \xi \leq x \leq \xi + \epsilon \\ 0 & \xi + \epsilon \leq x \leq b \end{cases} \quad (1)$$

If we substitute this function in the above expression and let $\epsilon \rightarrow 0$, then the term

$$\frac{1}{\epsilon^2} \int_{\xi-\epsilon}^{\xi+\epsilon} L_{y'y'} dx$$

will dominate the left hand side of the above inequality and determine its sign. Thus the sign of $L_{y'y'}$ determines the sign of $f''(0)$, and for minimum, the Legendre condition $L_{y'y'} \geq 0$ must hold. We have the Legendre test:

Theorem 1. *The Legendre test. If*

(i) Euler-Lagrange equation is satisfied,

(i) the range of integration is sufficiently small,
 (iii) the sign of $L_{y'y'}$ is constant throughout this range,
 then $J(y)$ is a minimum or a maximum value of J according as the sign of $L_{y'y'}$ is positive or negative.

The second variation: Let $y = y(x)$ be the path of integration for which the integral $J(y) = \int_a^b L(x, y, y')dx$ is minimum or maximum (sometimes called the integral is stationary). Consider all admissible functions $y + \varepsilon h$ where both y and h belong to $C^2[a, b]$. Then

$$J(y + \varepsilon h) = J(y) + \frac{\varepsilon^2}{2} J_2 + O(\varepsilon^3)$$

where

$$J_2 = \int_a^b [h^2 \frac{\partial^2 L}{\partial y^2} + 2h h' \frac{\partial^2 L}{\partial y \partial y'} + h'^2 \frac{\partial^2 L}{\partial y'^2}] dx$$

If $J(y)$ is minimum (or maximum) the sign of J_2 must be positive (or negative) without depending upon the choice of $h(x)$, for all sufficiently small ε . For simplicity we use the following notation: $L_0 = \frac{\partial L}{\partial y}$, $L_1 = \frac{\partial L}{\partial y'}$, $L_{00} = \frac{\partial^2 L}{\partial y^2}$, $L_{01} = \frac{\partial^2 L}{\partial y \partial y'}$, $L_{11} = \frac{\partial^2 L}{\partial y'^2}$. Then J_2 takes the form

$$J_2 = \int_a^b [L_{00} h^2 + 2L_{01} h h' + L_{11} h'^2] dx$$

We have the following Lemma. Proof is straightforward (done in the class)

Lemma 2. *Let $h(a) = h(b) = 0$, then*

$$J_2 = \int_a^b \{h^2 (L_{00} - \frac{d}{dx} L_{01}) - h \frac{d}{dx} (h' L_{11})\} dx$$

Definition. (*Jacobi equation and Jacobi field*). *Let $u \in C^2[a, b]$. Then the following equation is called the Jacobi equation.*

$$\{L_{00} - \frac{d}{dx}(L_{01})\} u - \frac{d}{dx}(L_{11} \frac{du}{dx}) = 0$$

the function u satisfying this second order ODE is called the Jacobi field

Lemma 3. If $h(a) = h(b) = 0$ and is a Jacobi field then

$$J_2 = \int_a^b L_{11} [h' - h \frac{u'}{u}]^2 dx,$$

If the term $\{h' - h(\frac{u'}{u})\} \neq 0$ and L_{11} has constant sign for all points of the extremal arc AB , where $A = (a, y(a))$ and $B = (b, y(b))$, then J_2 must have a sign which is independent of the choice of h . Now in the extremal case

$$J(y + \varepsilon h) - J(y) = \frac{\varepsilon^2}{2} J_2 + O(\varepsilon^3)$$

Hence J takes its minimum value (or maximum value) if L_{11} is positive (or negative) at all points on the extremal curve $y(x)$ with $x \in [a, b]$. This is essentially the Legendre test stated above.

If, however, L_{11} , does not keep its sign constant at all $x \in [a, b]$ on the curve $y(x)$, then the value $J(y)$ is neither a minimum nor a maximum.

If the term $\{h' - h(\frac{u'}{u})\}$ vanishes at all points of the extremal curve $y(x)$. Then it is clear that $h(x) = \alpha u(x)$, where α is an arbitrary constant. Along the extremal curve $y(x)$ the first variation vanishes. If in addition $h(x) = \alpha u(x)$ is chosen then the second variation, J_2 vanishes as well. The sign of $\delta J = J(y + \varepsilon h) - J(y)$ will depend on the third variation $\delta J = \frac{\varepsilon^3}{6} J_3$, where

$$J_3 = \int_a^b [h^3 L_{000} + 3h^2 h' L_{001} + 3hh'^2 L_{011} + h'^3 L_{111}] dx$$

Since the sign of δJ depends on that of ε there can be no maximum or minimum value of J unless J_3 vanishes, in which case the sign of δJ will depend on that of J_4 , the fourth variation. In order to avoid this difficulty

Jacobi proposed a test. This test provides whether the $h(x) = \alpha u(x)$ at all points of the extremal curve $y(x)$.

Definition: Let $u(x)$ be a solution of the Jacobi equation. Let $u(a) = 0$. This means that the Jacobi field vanishes at the point $A = (a, y(a))$. The all other points on the extremal curve $y(x)$ at which $u(x)$ vanishes are called the *conjugate points* to the point A .

The jacobi field (a function satisfying the Jacobi equation with $u(a) = 0$) may be given as the linear sum of the fundamental solutions u_1 and u_2 of the Jacobi equation.

$$u(x) = a_1 u_1(x) + a_2 u_2(x)$$

where a_1 and a_2 are constants. Since $u(a) = 0$, then

$$\frac{u_1(a)}{u_2(a)} = -\frac{a_1}{a_2}$$

Hence if x is the abscissa of the conjugate point to A ($u(x) = 0$) then

$$\frac{u_1(x)}{u_2(x)} = -\frac{a_1}{a_2}$$

It shows that this ratio is the same for all conjugate points. This is the way obtaining the conjugate points. We now give a nice way of determining the fundamental solutions u_1 and u_2 . Since the function $y(x)$, the solution of the Euler-Lagrange equation is second order ODE then it contains two independent constants of integration $y(x, c_1, c_2)$. It is easy to prove the following Lemma.

Lemma 4. $u_i = \frac{\partial y}{\partial c_i}$, $i = 1, 2$.

Hence the conjugate points are easily determined from the following formula.

Let $Y = (\frac{\partial y}{\partial c_1}) / (\frac{\partial y}{\partial c_2})$ then

$$Y(x) = Y(a)$$

Let A' be the first conjugate point to A when moving on the extremal curve $y(x)$ in the direction from $A = (a, y(a))$ to $B = (b, y(b))$. There are three possibilities

- (i) B lies between A and A' ,
- (ii) B coincides with A' ,
- (iii) B lies beyond A' .

In all three cases $u(x)$ vanishes at A , but in case (i) it can not vanish again at the point B , in case (ii) it vanishes again at B , and in case (iii) it vanishes again at some point of the arc AB lying between A and B . In case (i) we have $u(a) = h(a) = 0$ but $u(b) \neq 0$, $h(b) = 0$. Hence $u(x)$ can not be proportional to $h(x)$ at all points of the extremal arc AB . Therefore we must have $(h' - h\frac{u'}{u})^2 > 0$ at all points of AB , except, possibly, at finite number of points where $h(x)$ and $h'(x)$ vanish simultaneously. Then we have the following theorem *Jacobi test*.

Theorem 5. *Let $y = y(x)$ be the curve connecting the points A and B for which the functional $J(y) = \int_a^b L(x, y, y')dx$ takes its extremum value. Let A and A' be the conjugate points on this curve. If (i) B lies between A and A' and (ii) L_{11} has constant sign for all points on the arc AB , then $J(y)$ is a minimum (maximum) when $L_{11} > 0$ ($L_{11} < 0$)*

For all the other cases we need higher variations for extremum tests.