

CONSTRUCTING HOMOLOGICALLY TRIVIAL ACTIONS ON PRODUCTS OF SPHERES

ÖZGÜN ÜNLÜ AND ERGÜN YALÇIN

ABSTRACT. We prove that if a finite group G has a representation with fixity f , then it acts freely and homologically trivially on a finite CW-complex homotopy equivalent to a product of $f + 1$ spheres. This shows, in particular, that every finite group acts freely and homologically trivially on some finite CW-complex homotopy equivalent to a product of spheres.

1. INTRODUCTION

It is known that every finite group acts freely on a product of spheres $\mathbb{S}^{n_1} \times \cdots \times \mathbb{S}^{n_k}$ for some n_1, n_2, \dots, n_k . This follows from a construction given in [15, pg. 547] which is attributed to J. Tornehave by B. Oliver. The construction is based on a simple idea that one can permute the spheres in a product to get smaller isotropy. More specifically, for a finite group G , one defines the G -space X as a product of G -spaces $\text{Map}_{\langle g \rangle}(G, \mathbb{S}(\rho_g))$ over all elements $g \in G$, where $\mathbb{S}(\rho_g)$ denotes the unit sphere of a nontrivial one-dimensional complex representation $\rho_g : \langle g \rangle \rightarrow \mathbb{C}$. Note that because of the way X is constructed, G acts freely on X , but the induced action of G on the homology of X is not a trivial action in general. It is interesting to ask if there exists a similar construction so that the induced action on homology is trivial. The following was stated as a problem by J. Davis in the Problem Session of Banff 2005 conference on Homotopy Theory and Group Actions:

Question 1.1. Does every finite group act freely on some product of spheres with trivial action on homology?

To find a free, homologically trivial action of a finite group G on a product of spheres, one may try to take a family of G -spheres \mathbb{S}^{n_i} for $i = 1, \dots, k$ and let G act on the product $\mathbb{S}^{n_1} \times \cdots \times \mathbb{S}^{n_k}$ by diagonal action. An action which is obtained in this way is called a *product action* in general and a *linear action* if each G -sphere in the product is a unit sphere of a representation. By construction, product actions are homologically trivial if the action on each sphere is homologically trivial but not every finite group has a free product action on a product of spheres. For example, it is known that the alternating group A_4 cannot act freely on a product of spheres by a product action. This follows from a result of Oliver [15, Theorem 1] which says that A_4 does not act freely on any product of equal dimensional spheres with trivial action on homology. On the other hand, it can

Date: April 17, 2013.

2010 Mathematics Subject Classification. Primary: 57S25; Secondary: 55R91.

Both of the authors are partially supported by TÜBİTAK-TBAG/110T712.

be shown that A_4 acts freely on $\mathbb{S}^2 \times \mathbb{S}^3$ with trivial action on homology. So, one cannot answer the above question affirmatively by considering only the product actions.

The situation with A_4 is not an exceptional case. For example, if one searches through the characters of a finite group G and tries to see when G has a family of characters χ_1, \dots, χ_k such that G acts freely on $\mathbb{S}(\chi_1) \times \dots \times \mathbb{S}(\chi_k)$, then one notices that not many groups have such a family of characters. In fact, Urmie Ray [16] showed that finite groups which have free linear actions are very rare: If a finite group G has a free linear action on a product of spheres, then all nonabelian simple sections of G are isomorphic to A_5 or A_6 .

In this paper we attack the problem stated above using some more recent construction methods that were developed to study the rank conjecture. In particular, we use some ideas from our earlier papers [19] and [20] where we constructed free actions on products of spheres for finite groups which have representations with small fixity and for p -groups with small rank. Fixity of a representation V of a finite group G over a field F is defined as the maximum of dimensions $\dim_F V^g$ over all elements $g \in G - \{1\}$. If $\rho : G \rightarrow U(n)$ is a faithful complex representation with fixity f , then G acts freely on the space $X = U(n)/U(n - f - 1)$. For small values of f , one can modify the space X to obtain a free action on a product of $f + 1$ spheres (see [1] and [19]). In this paper we improve this result to all values of fixity for actions on finite CW-complexes homotopy equivalent to product of spheres.

Theorem 1.2. *Let G be a finite group. If G has a faithful complex representation with fixity f , then G acts freely on a finite complex X homotopy equivalent to a product of $f + 1$ spheres with trivial action on homology.*

We prove this theorem using a recursive method for constructing free actions. In fact in the construction we use symplectic representations instead of complex representations and obtain free actions on a products of spheres with dimensions $n_i \geq 3$. The recursive method involves the construction of a G -equivariant spherical fibration $p : E \rightarrow X$ over a given finite G -CW-complex X in each step. We require that the total space E is also homotopy equivalent to a finite G -CW-complex and that the G -action on E has smaller isotropy than the G -action on X . Once they are constructed, by taking fiber joins these fibrations can be replaced by G -fibrations which are non-equivariantly homotopy equivalent to trivial fibrations. This gives a G -action on a finite CW-complex Y homotopy equivalent to $X \times \mathbb{S}^N$ for some $N > 0$. Using this method, after some steps, one gets a free action on a product of spheres. This method was first developed by Connolly and Prassidis [4] and later used also in [2] and [18].

Since every finite group has a faithful complex representation, every finite group has a complex representation with fixity f for some positive integer f . Hence, as a corollary of Theorem 1.2 we obtain an affirmative answer to Question 1.1 in homotopy category.

Corollary 1.3. *Every finite group acts freely and homologically trivially on some finite CW-complex X homotopy equivalent to a product of spheres*

The paper is organized as follows: In Section 2, we introduce G -fibrations and discuss the effects of taking fiber joins of G -fibrations. In Section 3, we discuss the equivariant

Federer spectral sequence introduced by Møller [14] and using it, we give another proof for a theorem by M. Klaus [12] (see Theorem 3.1). In Section 4, we introduce our main construction method, and finally in Section 5, we prove Theorem 1.2.

2. G -FIBRATIONS

In this section, we first give some preliminaries on G -fibrations and then prove some lemmas on fiber joins of G -fibrations. For more details on this material we refer the reader to [13] and [22]. Some of this material also appears in [4], [9], [12], and [18].

Definition 2.1. A G -fibration is a G -map $p : E \rightarrow B$ which satisfies the following homotopy lifting property for every G -space X : Given a commuting diagram of G -maps

$$\begin{array}{ccc} X \times \{0\} & \xrightarrow{h} & E \\ \downarrow & & \downarrow p \\ X \times I & \xrightarrow{H} & B, \end{array}$$

there exists a G -map $\tilde{H} : X \times I \rightarrow E$ such that $\tilde{H}|_{X \times \{0\}} = h$ and $p \circ \tilde{H} = H$.

Given a G -fibration $p : E \rightarrow B$ over B , the isotropy subgroup $G_b \leq G$ of a point $b \in B$ acts on the fiber space $F_b := p^{-1}(b)$. So, F_b is a G_b -space. Let us denote the set of isotropy subgroups of the G -action on B by $\text{Iso}(B)$.

Definition 2.2. Let $\{F_H\}$ denote a family of H -spaces over all $H \in \text{Iso}(B)$. If for every $b \in B$, the fiber space F_b is G_b -homotopy equivalent to F_{G_b} , then $p : E \rightarrow B$ is said to have *fiber type* $\{F_H\}$.

Note that in general a G -fibration does not have to have a fiber type, i.e., for $b_1, b_2 \in B$ with $G_{b_1} = G_{b_2} = H$, it may happen that F_{b_1} and F_{b_2} are not H -homotopy equivalent. But throughout the paper we only consider G -fibrations which have a fiber type. Observe that if $p : E \rightarrow B$ is a G -fibration such that B^H is path connected for every $H \in \text{Iso}(B)$, then p has a fiber type since for every $b_1, b_2 \in B^H$, the fiber spaces F_{b_1} and F_{b_2} are H -homotopy equivalent by a standard argument in homotopy theory. In our applications the G -fibrations that we construct will often satisfy this connectedness property.

If $p : E \rightarrow B$ is a G -fibration with fiber type $\{F_H\}$ such that for all $H \in \text{Iso}(B)$ the fixed point space B^H is connected, then the family $\{F_H\}$ satisfies a certain compatibility condition. To see this, let $H, K \in \text{Iso}(B)$ such that $g^{-1}Kg \leq H$ for some $g \in G$. Then, we have $gB^H \subseteq B^K$, so by the connectedness of B^K , we obtain that for every $b \in B^H$, the K -space gF_b is K -homotopy equivalent to F_K . This means that the K -spaces $\text{Res}_K g^*F_H$ and F_K are K -homotopy equivalent for all $H \in \text{Iso}(B)$ where $\text{Res}_K g^*F_H$ is the space F_H which is considered as a K -space through the map $K \rightarrow H$ defined by $k \rightarrow g^{-1}kg$.

Definition 2.3. Let \mathcal{H} be a family of subgroups of G closed under conjugation. A family of H -spaces $\{F_H\}$ over all $H \in \mathcal{H}$ is called a *compatible family* of H -spaces if for every $H, K \in \mathcal{H}$ with $g^{-1}Kg \leq H$ for some $g \in G$, the K -space $\text{Res}_K g^*F_H$ is K -homotopy equivalent to F_K , where $\text{Res}_K g^*F_H$ is the the space F_H considered as a K -space through the map $K \rightarrow H$ defined by conjugation $k \rightarrow g^{-1}kg$.

The main aim of this section is to introduce some tools for construction of G -fibrations with fiber type $\{F_H\}$ for a given compatible family $\{F_H\}$. We first introduce some more terminology:

Given two G -fibrations $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ over the same G -space B , a G -map $f : E_1 \rightarrow E_2$ is called a fiber preserving map if it satisfies $p_2 \circ f = p_1$. Two fiber preserving G -maps $f, f' : E_1 \rightarrow E_2$ are said to be G -fiber homotopic if there is a G -map $H : E_1 \times I \rightarrow E_2$ which is fiber preserving at each $t \in I$ such that $H(x, 0) = f(x)$ and $H(x, 1) = f'(x)$ for all $x \in E_1$. We say two G -fibrations $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ are G -fiber homotopy equivalent if there are fiber preserving G -maps $f_1 : E_1 \rightarrow E_2$ and $f_2 : E_2 \rightarrow E_1$ such that $f_1 \circ f_2$ and $f_2 \circ f_1$ are G -fiber homotopic to identity maps.

For an H -space F_H , let $\text{Aut}_H(F_H)$ denote the topological monoid of self H -homotopy equivalences of F_H . Note that $\text{Aut}_H(F_H)$ is not a connected space in general but it is easy to show that all its components have the same homotopy type. When we need to choose a component, we often take the connected component which includes the identity map. We denote this component by $\text{Aut}_H^I(F_H)$.

Since $\text{Aut}_H(F_H)$ is a monoid, the usual construction of classifying spaces for monoids applies, and we get a universal fibration $E \text{Aut}_H(F_H) \rightarrow B \text{Aut}_H(F_H)$ with fiber $\text{Aut}_H(F_H)$. From this one also obtains a fibration

$$F_H \rightarrow E_H \rightarrow B \text{Aut}_H(F_H)$$

where $E_H = E \text{Aut}_H(F_H) \times_{\text{Aut}_H(F_H)} F_H$. This is actually an H -fibration with trivial H -action on the base space. It turns out that this fibration is a universal fibration for all H -equivariant fibrations with trivial action on the base space.

Theorem 2.4. *Let H be a finite group, F_H be a finite H -CW-complex, and B be a CW-complex with trivial H -action. Then, there is a one-to-one correspondence between H -fiber homotopy classes of H -fibrations over B with fiber F_H and the set of homotopy classes of maps $B \rightarrow B \text{Aut}_H(F_H)$. The correspondence is given by taking the pullback of the universal H -fibration described above via the map $f : B \rightarrow B \text{Aut}_H(F_H)$.*

This theorem is proved in [9] in full detail. The proof is based on the proof of Stasheff's theorem on the classification of Hurewicz fibrations [17]. More general versions of this theorem also appear in [8] and [22].

Also note that, as in the case of orientable vector bundle theory, we can give an orientable version of the classification of H -fibrations over a trivial H -space B . If $p : E \rightarrow B$ is an H -fibration with fiber F_H over a trivial H -space, then there is a natural group homomorphism $\chi : \pi_1(B) \rightarrow \pi_0(\text{Aut}_H(F_H)) = \mathcal{E}_H(F_H)$ where $\mathcal{E}_H(F_H)$ denotes the group of homotopy classes of self H -homotopy equivalences of F_H . If this homomorphism is trivial, then we call the H -fibration p a *homotopy orientable H -fibration*. This notion of orientability is stronger than usual notion of orientable fibration where one only requires the action on the homology of F_H to be trivial (see [6]). Note that an H -fibration is homotopy orientable if and only if its classifying map $f : B \rightarrow B \text{Aut}_H(F_H)$ lifts to a map $\tilde{f} : B \rightarrow B \text{Aut}_H^I(F_H)$. Also note that homotopy orientable fibrations are classified (as homotopy orientable fibrations) by the homotopy classes of maps $[B, B \text{Aut}_H^I(F_H)]$. We will be using these facts later in the proof of Lemma 2.8.

In the rest of this section we focus on the fiber join construction performed on G -fibrations. We use fiber joins to kill obstructions that occur in the construction of G -fibrations. The fiber join of two G -fibrations is defined in the following way: Let $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ be two fibrations. We define a fibration $E_1 \times_B E_2$ and maps $E_1 \times_B E_2 \rightarrow E_i$ for $i = 1, 2$ by the following pullback diagram

$$\begin{array}{ccc} E_1 \times_B E_2 & \longrightarrow & E_1 \\ \downarrow & & \downarrow p_1 \\ E_2 & \xrightarrow{p_2} & B. \end{array}$$

Then the G -space $E_1 *_B E_2$ is defined as the homotopy pushout of the following diagram

$$\begin{array}{ccc} E_1 \times_B E_2 & \longrightarrow & E_1 \\ \downarrow & & \downarrow \\ E_2 & \longrightarrow & E_1 *_B E_2. \end{array}$$

By the universal property of homotopy pushouts we get a G -fibration

$$p_1 * p_2 : E_1 *_B E_2 \rightarrow B$$

called the *fiber join of p_1 and p_2* . To see that $p_1 * p_2$ is a fibration, we can use equivariant lifting functions. Recall that a G -map $p : E \rightarrow B$ is a G -fibration if and only if there is a G -map

$$\lambda : \Omega_p = \{(e, \omega) \in E \times B^I \mid p(e) = \omega(0)\} \rightarrow E^I$$

such that $\lambda(e, \omega)(0) = e$ and $p\lambda(e, \omega) = \omega$ (see [23, pg. 30]). The G -map λ is called a *G -lifting function*. Let λ_1 and λ_2 denote the G -lifting functions for p_1 and p_2 . Then we can define a lifting function for the join $p_1 * p_2$ as follows

$$\lambda(te_1 + (1-t)e_2, \omega)(u) = t\lambda_1(e_1, \omega)(u) + (1-t)\lambda_2(e_2, \omega)(u)$$

for all t, u in I . So, $p_1 * p_2$ is a G -fibration.

Iterating this construction, we obtain a G -fibration

$$*_k p : \underbrace{E *_B E *_B \cdots *_B E}_{k\text{-many}} \rightarrow B$$

which we call the *k -fold fiber join of p with itself*. Note that if p is a G -fibration with fiber type $\{F_H\}$, then the fiber type of the k -fold join $*_k p$ is $\{*_k F_H\}$. If B has trivial H -action, then the k -fold join is classified by a map $B \rightarrow B \operatorname{Aut}_H(*_k F_H)$. We would like to explain this map in terms of the classifying map of p . For this, observe that there is a monoid homomorphism

$$\varphi : \operatorname{Aut}_H(F_H) \times \cdots \times \operatorname{Aut}_H(F_H) \rightarrow \operatorname{Aut}_H(*_k F_H)$$

defined by

$$\varphi(a_1, \dots, a_k)(x_1 t_1, \dots, x_k t_k) = (a_1(x_1)t_1, \dots, a_k(x_k)t_k)$$

for every $x_1, \dots, x_k \in F_H$ and $t_1, \dots, t_k \in [0, 1]$ with $\sum_i t_i = 1$. We have the following lemma:

Lemma 2.5. *Let H be a finite group, F_H be a finite H -CW-complex, and B be a CW-complex with trivial H -action. If $p : E \rightarrow B$ is a H -fibration with fiber type F_H whose classifying map is $f : B \rightarrow B \operatorname{Aut}_H(F_H)$, then the classifying map of the G -fibration $*_k p$ is given by the composition*

$$B \xrightarrow{f \times \cdots \times f} B \operatorname{Aut}_H(F_H) \times \cdots \times B \operatorname{Aut}_H(F_H) \xrightarrow{B\varphi} B \operatorname{Aut}_H(*_k F_H)$$

where $B\varphi$ is the map induced from the monoid homomorphism φ defined above.

Proof. Let $A = \prod_{i=1}^k \operatorname{Aut}_H(F_H)$. By standard properties of homotopy pushout diagrams, we observe that the fibration $*_k p$ is the pullback fibration of the fibration

$$q : EA \times_A (*_k F_H) \rightarrow BA$$

via the map $\prod_i f : B \rightarrow BA$. Note that A acts on $*_k F_H$ via the monoid homomorphism φ , so the classifying map of q is $B\varphi$. This completes the proof. \square

A special case of a G -fibration is a G -fiber bundle over a G -CW-complex. More specifically, if $\xi : E \rightarrow B$ is a complex G -vector bundle over a G -CW-complex B , then the sphere bundle $p : S(E) \rightarrow B$ of this vector bundle is a spherical G -fibration. Note that for every $b \in B$, the fiber space $p^{-1}(b)$ is a G_b -space which is homeomorphic to $\mathbb{S}(V_{G_b})$ where V_{G_b} denotes vector space $\xi^{-1}(b)$ with the induced G_b -action.

Note that when B^H is path connected for all $H \in \operatorname{Iso}(B)$, the family of complex representations $\{V_H\}$ defined over all $H \in \operatorname{Iso}(B)$ is a compatible family. The compatibility of a family of representations is defined in the following way: A family of representations $\alpha_H : H \rightarrow U(n)$ over $H \in \mathcal{H}$ is called a *compatible family of representations* if for every map $c_g : H \rightarrow K$ induced by conjugation with $g \in G$, there exists a $\gamma \in U(n)$ such that the following diagram commutes

$$\begin{array}{ccc} K & \xrightarrow{\alpha_K} & U(n) \\ \downarrow c_g & & \downarrow c_\gamma \\ H & \xrightarrow{\alpha_H} & U(n). \end{array}$$

Note that if F_H is an H -space which is H -homotopy equivalent to $\mathbb{S}(V_H)$ for some compatible family of complex H -representation V_H , then $\{F_H\}$ is a compatible family of H -spaces. So, the sphere bundle $p : S(E) \rightarrow B$ of a G -vector bundle is a spherical G -fibration with fiber type $\{\mathbb{S}(V_H)\}$. Note also that for every $k \geq 1$, the fiber join $*_k \mathbb{S}(V_H)$ is H -homotopy equivalent to the H -space $\mathbb{S}(V_H^{\oplus k})$ where $V_H^{\oplus k}$ denotes the k -fold direct sum of V_H .

In Section 4, we construct G -fibrations with fiber types of the form $\{\mathbb{S}(V_H)\}$. The following result is used in those constructions.

Lemma 2.6. *Let H be a finite group, F_H be an H -space which is H -homotopy equivalent to $\mathbb{S}(V_H)$ for some complex H -representation V_H . Let $\gamma, \gamma^1 : \operatorname{Aut}_H(F_H) \rightarrow \operatorname{Aut}_H(*_k F_H)$ be maps defined by $\gamma(a) = \varphi(a, a, \dots, a)$ and $\gamma^1(a) = \varphi(a, \operatorname{id}, \dots, \operatorname{id})$, respectively. Then, the induced group homomorphisms γ_* and γ_*^1 on homotopy $\pi_q(\operatorname{Aut}_H(F_H)) \rightarrow \pi_q(\operatorname{Aut}_H(*_k F_H))$ satisfy the relation $\gamma_* = k\gamma_*^1$.*

Proof. Let $\gamma^i : \text{Aut}_H(F_H) \rightarrow \text{Aut}_H(*_k F_H)$ be the map defined by

$$\gamma^i(a) = \varphi(\text{id}, \dots, a, \dots, \text{id})$$

where a is on the i -th coordinate. We have $\gamma = \gamma^1 \gamma^2 \dots \gamma^k$ under the product induced by the product in the monoid A . Since the group operation on $\pi_q(\text{Aut}_H(*_k F_H))$ coming from the monoid structure on $\text{Aut}_H(*_k F_H)$ coincides with the usual group structure on homotopy groups, we have $\gamma_* = \gamma_*^1 + \dots + \gamma_*^k$. So, to complete the proof, it is enough to show that γ^i and γ^j are homotopic for every $i, j \in \{1, \dots, k\}$. Since F_H is H -homotopy equivalent to $\mathbb{S}(V_H)$, it is enough to prove this for $\mathbb{S}(V_H)$. Note that in this case we have $\gamma^i = T(i, j) \gamma^j$ where $T(i, j) : V_H^{\oplus k} \rightarrow V_H^{\oplus k}$ is a linear transformation which swaps the j -th summand with the i -th summand. Since $U(n)$ is connected, there is a path between $T(i, j)$ and the identity. Using this path, we can define a homotopy between γ^i and γ^j . \square

Remark 2.7. For more general H -spaces F_H , there exists a swap map

$$S(i, j) : *_k F_H \rightarrow *_k F_H,$$

which swaps the i -th and j -th coordinates, similar to the linear transformation $T(i, j)$ in the proof of Lemma 2.6. If F_H is a free H -space homotopy equivalent to an odd dimensional sphere, then $S(i, j)$ will be homotopy equivalent to the identity map. If the H -action on F_H is not free, then the swap map $S(i, j)$ is not homotopy equivalent to the identity in general even when F_H is homotopy equivalent to an odd dimensional sphere. On the other hand, if F_H is a homotopy representation with the property that all fixed point spheres are odd dimensional, then under certain conditions on H or on the dimension function of F_H , one can prove that $S(i, j)$ is homotopy equivalent to the identity (see Proposition 20.12 in [13]).

We end this section with the following observation.

Lemma 2.8. *Let H be a finite group and $p : E \rightarrow \mathbb{S}^n$ be an H -fibration over the trivial H -space \mathbb{S}^n where $n \geq 2$. Suppose that the fiber type F_H of p is H -homotopy equivalent to $\mathbb{S}(V_H)$ for some complex H -representation V_H . If $\pi_{n-1}(\text{Aut}_H(F_H))$ is a finite group of order N , then $*_N p$ is H -fiber homotopy equivalent to the trivial fibration.*

Proof. By Theorem 2.4, the H -fibration p is classified by the homotopy class of a map $f : \mathbb{S}^n \rightarrow B \text{Aut}_H(F_H)$. Since $n \geq 2$, this map lifts to map $\tilde{f} : \mathbb{S}^n \rightarrow B \text{Aut}_H^I(F_H)$, so we can assume that the H -fibration p is an homotopy orientable fibration. Since $B \text{Aut}_H^I(F_H)$ is simply connected, we have

$$[\mathbb{S}^n, B \text{Aut}_H^I(F_H)] \cong \pi_n(B \text{Aut}_H^I(F_H)) \cong \pi_{n-1}(\text{Aut}_H(F_H)).$$

So, p is classified by a homotopy class $\alpha \in \pi_{n-1}(\text{Aut}_H(F_H))$. By a slightly modified version of Lemma 2.5, it is easy to see that the fiber join $*_N p$ is classified by $\gamma_*(\alpha)$ where $\gamma : \text{Aut}_H(F_H) \rightarrow \text{Aut}_H(*_N F_H)$ is the map defined by $\gamma(a) = \varphi(a, \dots, a)$. By Lemma 2.6, we have

$$\gamma_*(\alpha) = N \gamma_*^1(\alpha) = \gamma_*^1(N\alpha) = 0.$$

So, $*_N p$ is H -fiber homotopy equivalent to the trivial fibration. \square

3. EQUIVARIANT FEDERER SPECTRAL SEQUENCE

The main purpose of this section is to prove the following theorem which is due to M. Klaus [12]. We give a different proof here using the equivariant Federer spectral sequence which was introduced by Møller in [14].

Theorem 3.1 (Klaus [12]). *Let G be a finite group and V be a complex representation of G . Then, for every $n > 0$, there is an $m \geq 1$ such that $\pi_n(\text{Aut}_G(\mathbb{S}(V^{\oplus k})))$ is finite for all $k \geq m$.*

Before the proof, we first introduce Møller's theorem (Theorem 5.3 in [14]) and discuss how the Bredon cohomology with coefficients in a G -equivariant local coefficients system can be calculated for the case we are interested in. For more details on Bredon cohomology, we refer the reader to [3] and [13].

Let (X, A) be a finite G -CW-complex, $p : E \rightarrow B$ be a G -fibration, and $u : X \rightarrow E$ be a G -equivariant map. Assume that for all $H \leq G$ and for all $b \in B$, the space $p^{-1}(b)^H$ is a path connected simple space, and let $\pi_q(\mathcal{F})$ be the G -equivariant local coefficient system on X , induced from the local coefficient system on B associated to the G -fibration p (see Example 2.2 in [14]). Let $F_u(X, A; E, B)^G$ denote the space of all equivariant maps $v : X \rightarrow E$ such that $v|_A = u|_A$ and $p \circ v = p \circ u$ with compact open topology.

Theorem 3.2 (Møller [14]). *There is a spectral sequence with E^2 -term*

$$E_{pq}^2 = H_G^{-p}(X, A; \pi_q(\mathcal{F}))$$

for $p+q \geq 0$ and $E_{pq}^2 = 0$ otherwise, converging to $\pi_{p+q}(F_u(X, A; E, B)^G, u)$ when $p+q > 0$.

The spectral sequence above is called the equivariant Federer spectral sequence since it is the equivariant version of a spectral sequence introduced by Federer [7]. We will be using this spectral sequence for the following special case: Let X be a finite G -CW-complex such that X^H is a path connected simple space for all $H \leq G$. Take $A = \emptyset$, $E = X$, $B = *$, $p : E \rightarrow B$ to be the constant map, and $u : X \rightarrow E$ to be the identity map. Then $F_{\text{id}}(X, \emptyset; X, *)$ will be homotopy equivalent to the identity component of $\text{Aut}_G(X)$. Since all the components of $\text{Aut}_G(X)$ have the same homotopy type, we have

$$\pi_n(\text{Aut}_G(X)) \cong \pi_n(F_{\text{id}}(A, \emptyset; X, *))$$

for all $n > 0$. So, we can use the equivariant Federer spectral sequence to calculate the homotopy groups of $\text{Aut}_G(X)$.

Also note that in the situation we consider, the local coefficient system is constant on orbits. In this case Bredon cohomology with local coefficients has an alternative description which involves modules over the orbit category. We define this category now.

Definition 3.3. Let G be a finite group and Or_G denote the *orbit category* of G whose objects are orbits G/H where H is a subgroup of G . The morphisms of Or_G from G/H to G/K are G -maps $f : G/H \rightarrow G/K$, where we consider the left cosets G/H and G/K as left G -sets. We denote the morphism from G/H to G/K which sends H to aK by \hat{a} .

Let Γ denote the orbit category Or_G . A functor from Or_G to the category of abelian groups $\mathcal{A}b$ is called an $\mathbb{Z}\Gamma$ -module. Morphisms between $\mathbb{Z}\Gamma$ -modules are given by natural

transformations. Given a G -CW-complex X , we define a chain complex of $\mathbb{Z}\Gamma$ -modules by taking $C_n(X^?)$ as the functor $\text{Or}_G \rightarrow \mathcal{A}b$ which sends G/H to the n -th cellular chains $C_n(X^H)$ and sends $\hat{a} : G/H \rightarrow G/K$ to the group homomorphism $a_* : C_n(X^K) \rightarrow C_n(X^H)$. The boundary maps at each H are compatible with the maps a_* for every $\hat{a} : G/H \rightarrow G/K$, so they combine together to give a $\mathbb{Z}\Gamma$ -module map $\partial : C_n(X^?) \rightarrow C_{n-1}(X^?)$ for every n .

Given a $\mathbb{Z}\Gamma$ -module M , we define $C^*(G, M)$ as the cochain complex of abelian groups

$$C^n(X, M) = \text{Hom}_{\mathbb{Z}\Gamma}(C_n(X^?), M)$$

with coboundary maps coming from the boundary maps of $C_n(X^?)$. Note also that by the standard properties of free $\mathbb{Z}\Gamma$ -modules (see [13, Sec. 9]), we have

$$C^n(G, M) \cong \bigoplus_{[\sigma] \in \mathcal{I}_n} M(G/G_\sigma)$$

where \mathcal{I}_n is the set of G -orbits of n -cells in X .

Let \underline{L} be a G -equivariant local coefficient system on X (see [14, def. 2.1]). Suppose that there exists a $\mathbb{Z}\Gamma$ -module M such that $L(G/H)(x) = M(G/H)$ and $L(G/H)(\gamma) = \text{id}_{M(G/H)}$ for all $x \in X^H$ and all paths γ in X^H . Then we have

$$H_G^n(X; \underline{L}) \cong H^n(C^*(X, M))$$

where $H^n(G, \underline{L})$ denotes the Bredon cohomology with local coefficients and $C^*(G, M)$ denotes the cochain complex introduced above (see Definition 2.3 and Example 2.4 in [14]). In our situation, this gives an isomorphism

$$H_G^n(X; \pi_q(\mathcal{F})) \cong H_G^n(X; \pi_q(X^?))$$

where $\pi_q(X^?)$ is the $\mathbb{Z}\Gamma$ -module $\pi_q(X^?) : \text{Or}_G \rightarrow \mathcal{A}b$ which sends G/H to $\pi_q(X^H)$ and sends $\hat{a} : G/H \rightarrow G/K$ to $a_* : \pi_q(X^K) \rightarrow \pi_q(X^H)$. Also note that on the cochain level, we have

$$C^n(X, \pi_q(X^?)) = \text{Hom}_{\mathbb{Z}\Gamma}(C_n(X^?), \pi_q(X^?)) \cong \bigoplus_{[\sigma] \in \mathcal{I}_n} \pi_q(X^{G_\sigma}).$$

So, we have an explicit description of the E_{pq}^2 -terms of the equivariant Federer spectral sequence. Now we are ready to prove the main theorem of this section.

Proof of Theorem 3.1. Let G be a finite group and X be a finite G -CW-complex which is G -homotopy equivalent to $\mathbb{S}(V)$ for some complex representation V of G . In fact, we only need X to be a G -homotopy representation with odd dimensional fixed point spheres for our arguments to work (see [13, pg. 392] for a definition of homotopy representation). Let n be a fixed positive integer. We want to show that there is an $m \geq 1$ such that $\pi_n(\text{Aut}_G(*_k X))$ is finite for all $k \geq m$. Let $X_k = *_k X$ denote the k -fold join of X . By Theorem 3.2, there is a spectral sequence with

$$E_{pq}^2 = H_G^{-p}(X_k; \pi_q(X_k^?))$$

for $p + q \geq 0$ and $E_{pq}^2 = 0$ otherwise, converging to $\pi_{p+q}(\text{Aut}_G(X_k))$ when $p + q > 0$. Since X_k is finite dimensional, to show that $\pi_n(\text{Aut}_G(X_k))$ is finite it is enough to show that $H_G^{-p}(X_k; \pi_q(X_k^?))$ is finite for every pair (p, q) with $p + q = n$. Note that for this

we need to show that there is an $m \geq 1$ such that for all $k \geq m$, the cohomology group $H_G^{q-n}(X_k; \pi_q(X_k^?) \otimes \mathbb{Q})$ is zero for all $q \geq n$.

Let $\{n_1, n_2, \dots, n_s\}$ be the set of all distinct dimensions of fixed subspaces V^H over all subgroups $H \leq G$. Assume that $n_1 < n_2 < \dots < n_s$. Note that the fixed point spheres X_k^H have dimensions $\{kn_i - 1 \mid i = 1, \dots, s\}$. Since homotopy groups $\pi_i(S^{2j-1})$ of an odd dimensional sphere are all finite except when $i = 2j - 1$, we have $\pi_q(X_k^?) \otimes \mathbb{Q} = 0$ for all q which is not equal to $kn_i - 1$ for some i . If $q = kn_i - 1$ for some i , then we have

$$H_G^{q-n}(X_k; \pi_q(X_k^?) \otimes \mathbb{Q}) = H_G^{kn_i-n-1}(X_k; M_i)$$

where M_i is the $\mathbb{Z}\Gamma$ -module such that $M_i(H) = \mathbb{Q}$ for all subgroups $H \leq G$ satisfying $\dim V^H = n_i$. To complete the proof we need to show that this cohomology group is zero for all $i \in \{1, \dots, s\}$.

Note that there is a well-known first quadrant spectral sequence with E_2 -term

$$E_2^{pq} = \text{Ext}_{\mathbb{Z}\Gamma}^p(H_q(X_k^?), M_i)$$

which converges to $H_G^{p+q}(X_k; M_i)$ (see [20, Prop. 3.3]). Since the coefficient module M_i takes only the values \mathbb{Q} , we can replace $H_p(X_k^?)$ with $H_p(X_k^?; \mathbb{Q})$ and take the ext-groups over $\mathbb{Q}\Gamma$. Note that the $\mathbb{Q}\Gamma$ -module $H_p(X_k^?; \mathbb{Q})$ is zero at all dimensions except when $p = kn_i - 1$ for some i . Let N_i denote the $\mathbb{Q}\Gamma$ -module $H_{kn_i-1}(X_k^?; \mathbb{Q})$ for all $i = 1, \dots, s$. To prove that $H_G^{kn_i-n-1}(X_k; M_i) = 0$ for all i , it is enough to show that the ext-group

$$\text{Ext}_{\mathbb{Q}\Gamma}^{k(n_i-n_j)-n}(N_j, M_i)$$

is zero for all $j \leq i - 1$. Let l_j denote the length of the $\mathbb{Q}\Gamma$ -module N_j for every j (see [13, pg. 325] for a definition). Then, by [13, prop. 17.31], the above ext-group is zero if $k(n_i - n_j) - n \geq l_j$. Let $l = \max_j \{l_j\}$. Then if $k \geq n + l$, then the above inequality will hold for every $j \leq i - 1$. This completes the proof. \square

4. CONSTRUCTION OF SPHERICAL G -FIBRATIONS

We start with proving a proposition which is an important tool for constructing G -fibrations. In different forms, this proposition also appears in [4], [12], and [18]. Here we give a proof of it for completeness since it is the main ingredient in the proof of Theorem 1.2.

Proposition 4.1. *Let G be a finite group, B be a G -CW-complex, and let $\{V_H\}$ be a compatible family of complex representations over all $H \in \text{Iso}(B)$. Let $q_n : E_n \rightarrow B^{(n)}$, $n \geq 2$, be a G -fibration with fiber type $\{\mathbb{S}(V_H)\}$ where $B^{(n)}$ denotes the n -skeleton of B . Then there is an integer $k \geq 1$ and a G -fibration $q_{n+1} : E_{n+1} \rightarrow B^{(n+1)}$ such that the restriction of q_{n+1} to $B^{(n)}$ is G -fiber homotopy equivalent to $*_k q_n$. In particular, the fiber type of q_{n+1} is $\{\mathbb{S}(V_H^{\oplus k})\}$.*

Proof. By the definition of G -CW-complexes, there exists a pushout diagram

$$(1) \quad \begin{array}{ccc} \coprod_{i \in I_{n+1}} G/H_i \times \mathbb{S}^n & \xrightarrow{\coprod f_i} & B^{(n)} \\ \downarrow & & \downarrow \\ \coprod_{i \in I_{n+1}} G/H_i \times \mathbb{D}^{n+1} & \xrightarrow{\coprod g_i} & B^{(n+1)} \end{array}$$

where I_{n+1} is an indexing set of orbits of $(n+1)$ -cells in B . For each $i \in I_{n+1}$, let $q_{n,i}$ denote the G -fibration obtained by the following pullback diagram

$$\begin{array}{ccc} E_{n,i} & \longrightarrow & E_n \\ \downarrow q_{n,i} & & \downarrow q_n \\ G/H_i \times \mathbb{S}^n & \xrightarrow{f_i} & B^{(n)} \end{array} .$$

Restricting $q_{n,i}$ to the sphere \mathbb{S}^n in $G/H_i \times \mathbb{S}^n$ which is fixed by H_i , we obtain an H_i -fibration $q_{n,i}|_{\mathbb{S}^n} : q_{n,i}^{-1}(\mathbb{S}^n) \rightarrow \mathbb{S}^n$ such that the H_i -action on the base space is trivial. By Theorem 2.4 and by the argument in the proof of Lemma 2.8, such a fibration is classified by a homotopy class $\alpha_i \in \pi_{n-1}(\text{Aut}_{H_i}(\mathbb{S}(V_{H_i})))$.

By Theorem 3.1, for each $H \in \text{Iso}(B)$, there is an $m_H \geq 1$ such that $\pi_{n-1}(\text{Aut}_H(\mathbb{S}(V_H^{\oplus k})))$ is finite for all $k \geq m_H$. Let $m = \max\{m_H \mid H \in \text{Iso}(B)\}$. Then the group

$$\pi_{n-1}(\text{Aut}_H(\mathbb{S}(V_H^{\oplus m})))$$

has finite order, say d_H , for all $H \in \text{Iso}(B)$. Let $d = \prod_H d_H$. By Lemma 2.8, the H_i -fibration $*_{dm} q_{n,i}|_{\mathbb{S}^n}$ is H_i -fiber homotopy equivalent to the trivial fibration for all $i \in I_{n+1}$. This implies that the G -fibration p obtained by the following pullback diagram

$$\begin{array}{ccc} W & \xrightarrow{f} & *_{dm} E_n \\ \downarrow p & & \downarrow *_{dm} q_n \\ \coprod_{i \in I_{n+1}} G/H_i \times \mathbb{S}^n & \xrightarrow{\coprod f_i} & B^{(n)} \end{array}$$

is G -homotopy equivalent to the trivial fibration. Let

$$\varphi : \coprod_{i \in I_{n+1}} G \times_{H_i} \mathbb{S}(V_{H_i}^{\oplus dm}) \times \mathbb{S}^n \rightarrow W$$

be a G -fiber homotopy equivalence between the trivial fibration and p . We can use φ to glue the cone of the trivial fibration and obtain a G -quasifibration

$$q'_{n+1} : \left(\coprod_{i \in I_{n+1}} G \times_{H_i} \mathbb{S}(V_{H_i}^{\oplus dm}) \times \mathbb{D}^{n+1} \right) \cup_{f \circ \varphi} \left(*_{dm} E_n \right) \longrightarrow B^{(n+1)} .$$

A quasifibration $p : E \rightarrow B$ is a continuous map such that the induced map

$$p_* : \pi_i(E, p^{-1}(x), y) \rightarrow \pi_i(B, x)$$

is an isomorphism for all $x \in B$, $y \in p^{-1}(x)$, and $i \geq 0$ (see [5, def. 1.1]). A G -quasifibration is defined in an analogous way so that the corresponding modules over the orbit category are isomorphic (see [21, def. 2.1]). The fact that q'_{n+1} is a G -quasifibration follows from an equivariant version of Lemma 1.4 in [5] (see also [21, lem. 2.9]). There is a construction called gammafication that converts a quasifibration to a fibration and this construction also works for G -quasifibrations (see [21, Not. 2.8]). Applying gammafication to q'_{n+1} , we obtain a G -equivariant spherical fibration $q_{n+1} : E_{n+1} \rightarrow B^{(n+1)}$ whose fiber type is $\{\mathbb{S}(V_H^{\oplus dm})\}$. \square

Remark 4.2. Another possible way of completing the final step of the above construction is to attach trivial G -fibrations over $(n+1)$ -cells with the space $*_{dm}E_n$ using G -tubes (see [10, Theorem 3.1]). When these G -tubes are used one does not need the gammafication construction since one directly gets G -fibrations. This method is explained in detail in [9] and [10].

As a corollary of Proposition 4.1, we obtain the following which is also proved in [12] as Proposition 2.7.

Proposition 4.3 (Proposition 2.7, [12]). *Let G be a finite group and B be a finite dimensional G -CW-complex. Let $\{V_H\}$ be a compatible family of complex representations over all $H \in \text{Iso}(B)$. Then there exists an integer $k \geq 1$ and a G -equivariant spherical fibration $q : E \rightarrow B$ such that the fiber type of q is $\{\mathbb{S}(V_H^{\oplus k})\}$.*

Proof. Let

$$\mathbf{A} = (\rho_H) \in \lim_{\leftarrow H \in \mathcal{H}} \text{Rep}(H, U(n))$$

where $\mathcal{H} = \text{Iso}(B)$ and ρ_H is a representation for V_H for every $H \in \mathcal{H}$. Let $q : E_{\mathcal{H}}(G, \mathbf{A}) \rightarrow B_{\mathcal{H}}(G, \mathbf{A})$ denote the universal G -equivariant vector bundle with fiber type \mathbf{A} (see [20, def. 2.4]). Since $B_{\mathcal{H}}(G, \mathbf{A})^H = BC_{U(n)}(\rho_H)$ is simply connected for all $H \in \mathcal{H}$, by standard obstruction theory there is a G -map $B^{(2)} \rightarrow B_{\mathcal{H}}(G, \mathbf{A})$ (see the proof of Theorem 4.3 in [20] for details). Pulling back the universal G -equivariant bundle via this map, we obtain a G -equivariant vector bundle over $B^{(2)}$. The sphere bundle of this bundle is spherical G -fibration over $B^{(2)}$ with fiber type $\{\mathbb{S}(V_H)\}$. Now the result follows from the repeated application of Proposition 4.1. \square

We often want the total space of a G -fibration to be G -homotopy equivalent to a finite G -CW-complex. The following theorem gives a very useful criteria for this condition:

Proposition 4.4. *Let G be a finite group, B be a finite G -CW-complex, and $p : E \rightarrow B$ be a G -fibration with fiber type $\{F_H\}$. If F_H is H -homotopy equivalent to a finite H -CW-complex for every $H \in \text{Iso}(B)$, then E is G -homotopy equivalent to a finite G -CW-complex.*

Proof. We will prove this lemma by induction over the skeletons of B . We already know that $p^{-1}(B^{(0)})$ is G -homotopy equivalent to a finite G -CW-complex. Now assume that

$p^{-1}(B^{(n)})$ is G -homotopy equivalent to a finite G -CW-complex Z for some $n \geq 0$. We want to show that $p^{-1}(B^{(n+1)})$ is G -homotopy equivalent to a finite G -CW-complex.

The pushout diagram given in (1) induces a diagram of G -spaces

$$(2) \quad \begin{array}{ccc} f^*(E_n) & \xrightarrow{\bar{f}} & E_n \\ \downarrow \bar{j} & & \downarrow J \\ g^*(E_{n+1}) & \xrightarrow{\bar{g}} & E_{n+1} \end{array}$$

where the spaces in the diagram are the total spaces of the fibrations obtained by taking pullbacks of the fibration $q_{n+1} : E_{n+1} \rightarrow B^{(n+1)}$ via the maps f , g , j , and J . Here $f = \coprod f_i$, $g = \coprod g_i$,

$$j : \coprod_{i \in I_{n+1}} G/H_i \times \mathbb{S}^n \rightarrow \coprod_{i \in I_{n+1}} G/H_i \times \mathbb{D}^{n+1}$$

is the disjoint union of inclusion maps, and $J : B^{(n)} \rightarrow B^{(n+1)}$ is the inclusion map. Since the inclusion map $\mathbb{S}^n \rightarrow \mathbb{D}^{n+1}$ is a cofibration map, the G -map \bar{j} is a G -cofibration. So, by [13, Lemma 1.26], the diagram (2) is a pushout diagram and \bar{j} is a G -cofibration.

Since \mathbb{D}^{n+1} is contractible, there is a G -fiber homotopy equivalence

$$\coprod_{i \in I_{n+1}} G \times_{H_i} F_{H_i} \times \mathbb{D}^{n+1} \xrightarrow{\gamma} g^*(E_{n+1}).$$

This gives a commutative diagram of the following form

$$\begin{array}{ccccc} E_n & \xleftarrow{\bar{f}} & f^*(E_n) & \xrightarrow{\bar{j}} & g^*(E_{n+1}) \\ \uparrow \text{id} & & \uparrow \gamma' & & \uparrow \gamma \\ E_n & \xleftarrow{\bar{f} \circ \gamma'} & \coprod_{i \in I_{n+1}} G \times_{H_i} F_{H_i} \times \mathbb{S}^n & \xrightarrow{\text{id} \times j} & \coprod_{i \in I_{n+1}} G \times_{H_i} F_{H_i} \times \mathbb{D}^{n+1} \end{array}$$

where γ' is the restriction of γ to the boundary spheres. Such a restriction makes sense since γ is a fiber homotopy equivalence. Now, since both $\text{id} \times j$ and \bar{j} are G -cofibrations, by [13, Lemma 2.13], the G -space E_{n+1} , which is the pushout of the diagram in the first line, is G -homotopy equivalent to the pushout of the diagram in the second line.

To find a further homotopy equivalence, note that by induction assumption E_n is G -homotopy equivalent to a finite G -CW-complex Z . So, using a similar diagram as above, we can conclude that E_{n+1} is G -homotopy equivalent to the pushout of a diagram of the following form

$$Z \xleftarrow{\varphi} \coprod_{i \in I_{n+1}} G \times_{H_i} F_{H_i} \times \mathbb{S}^n \xrightarrow{\text{id} \times j} \coprod_{i \in I_{n+1}} G \times_{H_i} F_{H_i} \times \mathbb{D}^{n+1}.$$

Now, we can replace the map φ with a cellular one (up to homotopy) and conclude that E_{n+1} is G -homotopy equivalent to a finite G -CW-complex since the spaces Z and F_{H_i} for all $i \in I_{n+1}$ are finite G -CW-complexes. This completes the n -th stage of our induction.

Since B is a finite G -CW-complex, the induction will stop in finite steps. So, the proof is complete. \square

5. PROOF OF THE MAIN THEOREM

Now, we are ready to prove the main theorem of the paper. First we introduce some notation and recall some basic facts about Stiefel manifolds. For more details, we refer the reader to [19].

Let F denote the field of real numbers \mathbb{R} , complex numbers \mathbb{C} , or quaternions \mathbb{H} . For a real number the conjugation is defined by $\bar{x} = x$, for a complex number $x = a + ib$ by $\bar{x} = a - ib$, and for a quaternion $x = a + ib + jc + kd$ by $\bar{x} = a - ib - jc - kd$. On the vector space F^n , we can define an inner product (v, w) by taking

$$(v, w) = v_1\bar{w}_1 + v_2\bar{w}_2 + \cdots + v_n\bar{w}_n.$$

The Stiefel manifold $V_k(F^n)$ is defined as the subspace of F^{nk} formed by the k -tuples of vectors (v_1, v_2, \dots, v_k) such that $v_i \in F^n$ and for every pair (i, j) , we have $(v_i, v_j) = 1$ if $i = j$ and zero otherwise.

There is a sequence of fiber bundles

$$V_n(F^n) \rightarrow \cdots \rightarrow V_{k+1}(F^n) \xrightarrow{q_k} V_k(F^n) \rightarrow \cdots \rightarrow V_2(F^n) \xrightarrow{q_1} V_1(F^n)$$

where the map $q_k : V_{k+1}(F^n) \rightarrow V_k(F^n)$ is defined by $q_k(v_1, \dots, v_{k+1}) = (v_1, \dots, v_k)$ and the fiber of q_k is $V_1(F^{n-k}) = \mathbb{S}^{c(n-k)-1}$ where $c = \dim_{\mathbb{R}} F$ (see Theorem 3.8 and Corollary 3.9 in Chapter 8 of [11]). Note that the sphere bundle $q_k : V_{k+1}(F^n) \rightarrow V_k(F^n)$ is the sphere bundle of the vector bundle $\bar{q}_k : \bar{V}_{k+1}(F^n) \rightarrow V_k(F^n)$ where $\bar{V}_{k+1}(F^n)$ is the space formed by $(k+1)$ -tuples (v_1, \dots, v_{k+1}) satisfying $(v_1, \dots, v_k) \in V_k(F^n)$ and $(v_i, v_{k+1}) = 0$ for all $i = 1, \dots, k$.

Note that if a finite group G has a representation W over a field F , then the inner product above can be replaced by a G -invariant one and the Stiefel manifolds have natural G -actions. Moreover the sphere bundles given above become G -equivariant bundles. If the representation W has fixity f , then we have $\dim_F W^g \leq f$ for all $g \in G - \{1\}$. This means if W is a faithful representation, i.e., $f \leq \dim_F W - 1$, then the G -action on $V_{f+1}(W)$ is free. We will be using this observation in the proof of the main theorem.

Also observe that if G has a complex representation with fixity f , then by tensoring it with \mathbb{H} over \mathbb{C} , we obtain a symplectic representation with the same fixity. So to prove Theorem 1.2, it is enough to prove the same statement for symplectic representations. On the other hand with symplectic representations unit spheres of fixed point subspaces are simply connected, so it makes the proof easier. So, we prove the following:

Theorem 5.1. *Let $\rho : G \rightarrow Sp(n)$ be a faithful symplectic representation with fixity f . Then there exists a finite G -CW-complex X homotopy equivalent to a product of $f + 1$ spheres such that the G action the homology of X is trivial.*

Proof. Let W denote the \mathbb{H} -space corresponding to the representation ρ . Define $X_1 = V_1(W)$. We will construct finite G -CW-complexes X_2, X_3, \dots, X_{f+1} recursively. For all i , the G -CW-complex X_i will be homotopy equivalent to a product of i spheres and will satisfy the following property: If $H \in \text{Iso}(X_i)$, then $V_i(W)^H \neq \emptyset$ where $\text{Iso}(X_i)$ denotes the set of isotropy subgroups of X_i .

Assume that X_i is constructed for some $i \geq 1$. Note that for every $H \leq G$, the fixed point set $V_i(W)^H$ is either empty or simply connected. Since for every $H \in \text{Iso}(X_i)$ we have $V(W_i)^H \neq \emptyset$, by standard equivariant obstruction theory there exists a G -map $f : X_i^{(2)} \rightarrow V_i(W)$. By pulling back the G -bundle $q_i : V_{i+1}(W) \rightarrow V_i(W)$ via f , we obtain a G -equivariant vector bundle over $X_i^{(2)}$ with fiber type $\{S(V_H)\}$. Note that this is a compatible family defined over all $H \in \text{Iso}(X_i)$ where V_H is the H -space $(\overline{q}_i)^{-1}(b)$ for some $b \in V_i(W)^H$.

Now, applying Proposition 4.1 repeatedly to this G -fibration, we obtain a spherical G -fibration $E_i \rightarrow X_i$ with fiber type $\{S(V_H^{\oplus k})\}$ for some $k \geq 1$. By taking further fiber joins, we can assume that E_n is a trivial fibration non-equivariantly and the action on the homology of the total space is trivial. This is shown below in Lemma 5.2. Now, by Proposition 4.4, E_i is G -homotopy equivalent to a finite G -CW-complex Y . Hence we can take X_{i+1} as Y and continue the induction until we reach X_{f+1} . At this stage, we have $(V_{f+1}(W))^H \neq \emptyset$ implies $H = \{1\}$, so we can conclude that the G -action on X_{f+1} is free. \square

Lemma 5.2. *Let $p : E \rightarrow B$ be a G -fibration over a finite G -CW-complex B and n be a positive integer. Suppose that p has fiber type $\{F_H\}$ such that F_H is homotopy equivalent to the sphere \mathbb{S}^n for all $H \in \text{Iso}(B)$. Then, there is an integer $k \geq 1$ such that $*_k p : *_k E \rightarrow B$ is non-equivariantly homotopy equivalent to the trivial fibration. Moreover, if the G -action on the cohomology of B is trivial, then there is an integer $m > 0$ such that for every even number $k > m$, the G -action on the cohomology of $*_k E$ is also trivial.*

Proof. The first part of the lemma is well-known and it follows from the fact that the homotopy groups of $\text{Aut}(S^n)$ is finite. For the second part, observe that since the resulting fibration is homotopy equivalent to the trivial fibration, it is in particular an orientable fibration, i.e., $\pi_1(B)$ action on the homology of F is trivial. So, there exists a consistent choice of generators for $H^n(F_b) \cong \mathbb{Z}$ for all $b \in B$. Note that this gives a G -action on the cohomology of the fibers $H^n(F_b)$ which is defined by

$$g^* : H^n(F_b) \rightarrow H^n(F_{gb}) \cong H^n(F_b)$$

where the isomorphism on the right comes from identifications of generators that we have chosen. Observe that this action in general can be nontrivial since a generator u can go to $-u$ but if we take the fiber join of p with itself, then we can assume that this action is trivial for all $b \in B$. So for every even number k , the fibration $*_k p$ will have this property.

For an orientable spherical fibration there is a Serre spectral sequence

$$E_2^{p,q} = H^p(B, H^q(F)) \implies H^{p+q}(E).$$

Note that for a G -fibration, all the terms in this spectral sequence will be $\mathbb{Z}G$ -modules and the differentials will be $\mathbb{Z}G$ -module homomorphisms since every $g \in G$ induces a fiber preserving continuous map $g : E \rightarrow E$. In our case, we have a two line spectral sequence and it is easy to see that by choosing k large enough, we can assume that $N = \dim(*_k \mathbb{S}^n) \geq \dim B$, and hence we can conclude that $H^i(E) \cong H^i(B)$ for $i < N$ and $H^i(E) \cong H^{i-N}(B, H^N(F))$ for $i \geq N$. Since G acts trivially on $H^*(B)$, we have trivial action on $H^*(E)$ if the G -action on $H^{i-N}(B, H^N(F))$ is trivial for all $i \geq N$. Note that

the Serre spectral sequence has a product structure, so the action on $H^*(B, H^N(F))$ is trivial if the G -action on $H^0(B, H^N(F))$ is trivial. Note that $H^0(B, H^N(F))$ is the kernel of the map

$$d^1 : \text{Hom}_{\mathbb{Z}}(C_0(B), H^N(F)) \rightarrow \text{Hom}_{\mathbb{Z}}(C_1(B), H^N(F))$$

and as a $\mathbb{Z}G$ -module

$$\text{Hom}_{\mathbb{Z}}(C_i(B), H^N(F)) \cong \bigoplus_{\sigma \in I_i} \text{Hom}_{\mathbb{Z}}(\text{Ind}_{G_\sigma}^G \mathbb{Z}_\sigma, H^N(F_\sigma))$$

for $i = 0, 1$ where the G -action on $H^N(F_\sigma)$ is the one described above. So if k is a large even number, then this action is trivial, and we can conclude that $H^0(B, H^N(F)) \cong H^0(B)$ as $\mathbb{Z}G$ -modules, and hence $H^0(B, H^N(F))$ is a trivial $\mathbb{Z}G$ -module. This completes the proof of the lemma. \square

Acknowledgement. We thank Natalia Castellana for directing us towards Møller's work on equivariant Federer spectral sequence for calculating the homotopy groups of the monoid $\text{Aut}_H(F_H)$. We also thank the referee for careful reading of the paper and for corrections and remarks on the first version of the paper.

REFERENCES

- [1] A. Adem, J. F. Davis, and Ö. Ünlü, *Fixity and free group actions on products of spheres*, Comment. Math. Helv. **79** (2004), 758–778.
- [2] A. Adem and J. H. Smith, *Periodic complexes and group actions*, Ann. of Math. (2) **154** (2001), 407–435.
- [3] G. Bredon, *Equivariant Cohomology Theories*, Lecture Notes in Mathematics 34, Springer-Verlag, 1967.
- [4] F. Connolly and S. Prassidis, *Groups which act freely on $\mathbb{R}^m \times \mathbb{S}^{n-1}$* , Topology **28** (1989), 133–148.
- [5] A. Dold and R. Lashoff, *Principal quasi-fibrations and fibre homotopy equivalence of bundles*, Illinois J. Math. **3** (1959), 285–305.
- [6] K. Ehrlich, *The obstruction to the finiteness of the total space of a fibration*, Michigan Math. J. **28** (1981), 19–38.
- [7] H. Federer, *A study of function spaces by spectral sequences*, Trans. Amer. Math. Soc. **82** (1956), 340–361.
- [8] C. French, *The equivariant J-homomorphism*, Homology Homotopy Appl. **5** (2003), 161–212.
- [9] A. Güçlükan İlhan, *Obstructions for constructing G-equivariant fibrations*, PhD. thesis (2011).
- [10] A. Güçlükan İlhan, *Obstructions for constructing equivariant fibrations*, preprint, arXiv:1110.3880.
- [11] D. Husemoller, *Fibre Bundles*, Third edition, Graduate Texts in Mathematics 20, Springer-Verlag, New York, 1994.
- [12] M. Klaus, *Constructing free actions of p-groups on products of spheres*, Algebr. Geom. Topol. **11** (2011), 3065–3084.
- [13] W. Lück, *Transformation Groups and Algebraic K-Theory*, Lecture Notes in Mathematics, vol. 1408, Springer-Verlag, Berlin, 1989.
- [14] J. M. Møller, *On equivariant function spaces*, Pacific J. Math. **142** (1990), 103–119.
- [15] R. Oliver, *Free compact group actions on products of spheres*, Algebraic Topology: Aarhus, Denmark 1978, Lecture Notes in Mathematics 763, Springer-Verlag, Berlin, 1979, pp. 539–548.
- [16] U. Ray, *Free linear actions of finite groups on products of spheres*, J. Algebra **147** (1992), 456–490.
- [17] J. Stasheff, *A classification theorem for fiber spaces*, Topology **2** (1963), 239–246.
- [18] Ö. Ünlü, *Constructions of free group actions on products of spheres*, PhD. thesis (2004).

- [19] Ö. Ünlü and E. Yalçın, *Quasilinear actions on products of spheres*, Bull. London Math. Soc. **42** (2010), 981–990.
- [20] Ö. Ünlü and E. Yalçın, *Fusion systems and constructing free actions on products of spheres*, Math. Z. **270** (2012), 939–959.
- [21] S. Waner, *Equivariant fibrations and transfer*, Trans. Amer. Math. Soc. **258** (1980), 369–384.
- [22] S. Waner, *Equivariant classifying spaces and fibrations*, Trans. Amer. Math. Soc. **258** (1980), 385–405.
- [23] G. W. Whitehead, *Elements of Homotopy Theory*, Graduate Texts in Mathematics 61, Springer, 1978.

DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, ANKARA, 06800, TURKEY.
E-mail address: unluo@fen.bilkent.edu.tr, yalcine@fen.bilkent.edu.tr